

ชยาธิ: การกำจัดของเสียของครัวเรือน: ความร การรบรร การปฏิบัติ และความสัมพันธ์กับความถี่ของ การเกิดโรคอุจจาระร่วง ในเมืองลาพุดตา ประเทศพม่า (HOUSEHOLD WASTE DISPOSAL: KNOWLEDGE, PERCEPTION, PRACTICES, AND RELATIONSHIP WITH DIARRHEA FREQUENCY IN LAPUTTA TOWNSHIP IN MYANMAR) อ.ที่ปรึกษาวิทยานิพนธ์หลัก: Robert S. Chapman, M.D., M.P.H., 98 หน้า

การศึกษาแบบภาคตัดขวางนี้ได้ดำเนินการในหมู่บ้าน เฮ ทวิน เสก เมืองลาพุดตา ประเทศพม่า ในเดือนมีนาคมพ.ศ.2554โดยมีวัตถุประสงค์เพื่อประเมินระดับความรู้การรับรู้และการปฏิบัติในการกำจัดของเสียของครัวเรือนรวมทั้งเพื่อศึกษาความสัมพันธ์ระหว่างลักษณะทางประชากรสังคมลักษณะครัวเรือนระดับความรู้ การรับรู้และการปฏิบัติในการกำจัดของเสียของครัวเรือนกับความถี่ของการเกิดโรคอุจจาระร่วงในประชาชนเมือง ลาพุดตาประเทศพม่ากลุ่มตัวอย่างมีจำนวนทั้งสิ้น389คนซึ่งแบบสอบถามแบบมีโครงสร้างโดยผู้สัมภาษณ์เป็นผู้กรอกแบบสอบถามสถิติANOVAใช้ในการทดสอบความสัมพันธ์ระหว่างตัวแปรต้นได้แก่ลักษณะทางประชากร สังคมและลักษณะครัวเรือน กับการปฏิบัติในการกำจัดของเสียของครัวเรือน สำหรับการทดสอบ Chi-Square ใช้อธิบายความสัมพันธ์ระหว่างคุณลักษณะทางสังคมประชากรและครัวเรือนที่มีการเกิด โรคอุจจาระร่วงกับความ รู้และการรับรู้ในการกำจัดของเสียกับการปฏิบัติในการกำจัดของเสียของครัวเรือนเช่นเดียวกับความรู้และการรับรู้ ของการเกิดโรคอุจจาระร่วง

ผลการศึกษาในภาพรวมพบว่ามีการเกิดโรคอุจจาระร่วงเกิดขึ้น 19.5% โดยกลุ่มตัวอย่าง 78.1% มีความรู้ในระดับสูงและ21.1%อยู่ในระดับปานกลางสำหรับการรับรู้ในการกำจัดของเสียของครัวเรือนพบว่า 50.4%ของกลุ่มตัวอย่างมีการรับรู้ในระดับสูงและ48.1%อยู่ในระดับปานกลางการปฏิบัติในการกำจัดของเสียของครัวเรือนมีความแตกต่างกันอย่างมีนัยสำคัญเมื่อเปรียบเทียบแหล่งน้ำดื่ม ($p<0.001$) จำนวนถังขยะ ($p=0.032$) ถังขยะแบบมีฝาปิด ($p=0.01$) ถังขยะเปล่า ($p=0.008$) ขยะจากห้องครัว ($p=0.017$) และการรับรู้ต่อการกำจัดขยะในครัวเรือน($p<0.001$)กล่าวคือครัวเรือนที่ใช้แหล่งน้ำดื่มจากบ่อน้ำและน้ำฝนมีการปฏิบัติในการกำจัดของเสียครัวเรือนสูงครัวเรือนที่มีจำนวนถังขยะมากมีการปฏิบัติในการกำจัดของเสียครัวเรือนสูงครัวเรือนที่มีถังขยะแบบมีฝาปิดและการมีความถี่ในการทิ้งขยะมากมีการปฏิบัติในการกำจัดของเสียครัวเรือนสูง การใช้ส้วมในบ้าน (OR=2.38; 95% CI: 1.02-5.49) และจำนวนเด็กอายุต่ำกว่า 5 ปี (OR=2.09;95% CI: 1.08-4.04) ส่วนการใช้ถังขยะแบบมีฝาปิดสามารถป้องกัน โรคอุจจาระร่วงได้ (OR=0.30, 95% CI: 0.16-0.59)

การศึกษานี้เป็นการรวบรวมข้อมูลพื้นฐานเกี่ยวกับการกำจัดของเสียในครัวเรือนเพื่อประโยชน์สำหรับการศึกษาวิจัยต่อไปผลการศึกษานี้แสดงให้เห็นถึงความสำคัญในการกำหนดยุทธศาสตร์และการวางแผนของผู้นำกำหนดนโยบายเพื่อปรับปรุงวิธีการกำจัดของเสียของครัวเรือนและลดการเกิดโรคอุจจาระร่วงในเมืองลาพุดตา ประเทศพม่า

สาขาวิชา สาธารณสุขศาสตร์

ลายมือชื่อ นิลิต

ปีการศึกษา 2553

ลายมือชื่อ อ.ที่ปรึกษาวิทยานิพนธ์หลัก

##5379120453: MAJOR PUBLIC HEALTH

KEYWORDS: HOUSEHOLD WASTE DISPOSAL/ DIARRHEA FREQUENCY/ LAPUTTA TOWNSHIP/ MYANMAR

YE PAING KYI: HOUSEHOLD WASTE DISPOSAL: KNOWLEDGE, PERCEPTION, PRACTICES, AND RELATIONSHIP WITH DIARRHEA FREQUENCY IN LAPUTTA TOWNSHIP IN MYANMAR. ADVISOR: ROBERT S. CHAPMAN, M.D., M.P.H., 98pp.

A cross-sectional study was done in Yay Twin Seik Village Tract in Laputta Township, Myanmar, in March, 2011. The main objectives of this study were to assess the level of knowledge, perception and practices on household waste disposal; and to find out the association between socio-demographic, household characteristics, level of knowledge and perception with level of practices on household waste disposal associated with diarrhea frequency in Laputta residents. This study was carried out with 389 subjects by using a structured interviewer-administrated questionnaire to acquire data, with ethical review protocol no. 031.1/54 which was approved on 24th March, 2011. To find out the association between independent variables, socio-demographic and household characteristics; and practice on household waste disposal, the one-way analysis of variance (ANOVA) test was used. Chi-square tests were done to assess the associations between socio-demographic and household characteristics with diarrhea; knowledge and perception towards household waste disposal with practice on household waste disposal as well as with diarrhea occurrence.

The overall diarrhea occurrence was 19.5%. Among respondents, 78.1% had high level of knowledge while 21.1% had moderate level of knowledge. For perception towards household waste disposal, 50.4% of respondents had high-level perception and 48.1% had moderate level of perception. Practice on household waste disposal differed significantly with drinking water source (<0.001), number of trash bin ($p=0.032$), trash bins with wide lids ($p=0.01$), emptying of trash bins ($p=0.008$), kitchen waste ($p=0.017$) and perception towards household waste disposal (<0.001). The households which used pond water and rain water had higher practice score; the more the number of the trash bins, the higher the practice score; the use of trash bins with wide lids was linked to the higher practice score; and the more frequent they emptied trash bins, the more practice score they had. Indoor latrine (OR=2.38; 95% CI: 1.02-5.49) and number of children under 5 (OR=2.09; 95% CI: 1.08-4.04) were risk factors for diarrhea while trash bins with narrow lids (OR=0.30, 95% CI: 0.16-0.59) were protective against diarrhea.

This study provides baseline data regarding household waste disposal for further studies. The result can also inform policy makers to develop strategy and planning to improve household waste disposal and to reduce diarrhea occurrence. Further research on determinants of waste disposal practices and diarrhea occurrence is needed in Laputta and elsewhere.

Field of Study: ...Public Health.....Student's Signature.....

Academic Year: ..2010.....Advisor's Signature.....