
INCREASING THE IMMUNIZATION COVERAGE: AN INTERVENTION THROUGH THEMOBILIZATION OF MATERNAL AND CHILD HEALTH WORKERS IN NEPAL



© College of Public Health, Chulalongkorn University Bangkok, Thailand.

ABSTRACT

IMR of Nepal is very high (86 / 1000 live births) in comparison with other countries of the world. The main causes of the high IMR are vaccine preventable diseases, ARI, and diarrhea. It can be reduce by appropriate immunization to the under five children.

The main issue is that, how to increase the immunization coverage in Nepal. In developing countries this can only be achieve through the PHC approach. Community Health Workers are the backbone of this program. So, appropriate training regarding EPI and community participation, and effective supervision of Maternal and Child Health Workers will be helpful to improve the immunization coverage in Nepal.

A Seven days EPI training will be carried out in July 1998 by the researcher with the help of Regional Training Centre, Surkhet, and District Health Office, Dang. After the training, MCHWs will provide the immunization service to the community in the Dang district. Impact evaluation will be carried out after one year of implementations of program. The methods of data collection will be official statistics, FGDs and survey questionnaire. Results of study will be compared with control district (Bardiya), which is geographically and demographically similar to Dang district.

จุฬาลงกรณ์มหาวิทยาลัย
COLLEGE OF PUBLIC HEALTH SCIENCES
CHULALONGKORN UNIVERSITY